2026 Volume: 02 Issue: 11 www.ijrpa.com ISSN 2456-9995 Review Article

International Journal Research Publication Analysis

Page: 01-06

HAHNEMANN'S CONCEPTS OF MENTAL DISEASE AND THEIR CONCEPTUAL RELEVANCE TO PSYCHODERMATOLOGY

Dr Angela B. Williams*[1], Dr Manoj Narayan V.[2], Dr M. Murugan[3]

[1]Senior Resident, Dept of Organon of Medicine and Homoeopathic Philosophy, Sarada Krishna Homoeopathic Medical College and Hospital, (Affiliated to The Tamil Nadu Dr. M.G.R Medical University) Kulasekharam, Kanyakumari district, Tamil Nadu – 629161.

[2]Prof., Dept of Organon of Medicine and Homoeopathic Philosophy, Sarada Krishna Homoeopathic Medical College and Hospital, (Affiliated to The Tamil Nadu Dr. M.G.R. Medical University) Kulasekharam, Kanyakumari district, Tamil Nadu – 629161.

[3]PG &Ph.D. Guide, Department of Organon of Medicine, Sarada Krishna Homoeopathic Medical College and Hospital, (Affiliated to The Tamil Nadu Dr M.G.R. Medical University) Kulasekharam, Kanyakumari district, Tamil Nadu – 629161.

Article Received: 27 September 2026
Article Revised: 17 October 2026
Published on: 06 November 2026

***Corresponding Author: Dr Angela B. Williams**
Senior Resident, Dept of Organon of Medicine and Homoeopathic Philosophy, Sarada Krishna Homoeopathic Medical College and Hospital, (Affiliated to The Tamil Nadu Dr. M.G.R Medical University) Kulasekharam, Kanyakumari district, Tamil Nadu – 629161.

DOI: https://doi-doi.org/101555/ijrpa.7367

ABSTRACT

Psychodermatology explores the intricate relationship between psychological experiences and skin disorders, recognising the influence of emotional stress on cutaneous health and the psychological burden of chronic skin conditions. Hahnemann's concepts of mental diseases, particularly §§211–230 of the Organon of Medicine, emphasise individual disposition and the interrelationship between mental and physical manifestations. This article explores the conceptual relevance of these principles in understanding psychological affections in dermatological conditions. The discussion signifies the importance of emotional triggers, symptom chronology, psychological wellbeing, and comprehensive patient assessment. Although classical homoeopathic philosophy and modern psychoneuroimmunology represent distinct theoretical frameworks, their comparison offers an opportunity for interdisciplinary understanding of the mind–skin relationship. Integrating physical and psychological assessment may promote a more holistic, empathetic, and patient-centred approach to dermatological care.

KEYWORDS: Psychodermatology; Hahnemann; Mental Diseases; Mind–Skin Relationship; Organon of Medicine.

INTRODUCTION

Psychodermatology is the branch of interdisciplinary field of medicine that examines the interaction between psychological processes and skin disorders.[1] Skin diseases may be precipitated or aggravated by psychological stress, while chronic or visible skin conditions may contribute to anxiety, depression, low self-esteem, social withdrawal, and impaired quality of life.[2] These interactions are understood through the biopsychosocial model and the neuro-immuno-cutaneous system, which connects the nervous, endocrine, immune, and cutaneous systems.[3] Hahnemann discusses the importance of mental and emotional symptoms in understanding the individual patient, the relationship between physical and mental manifestations, and the need to consider the complete symptom picture. These concepts can be correlated with psychodermatology for studying the relationship between emotional experiences and cutaneous manifestations. The patient with a skin disease should be assessed not merely by the visible lesion but also by the associated emotional state, the chronology of symptoms, the influence of stress, and the impact of the disease on daily life. This correlation is useful for clinical observation and case analysis.[4] However, Hahnemann's theory of miasms and homoeopathic treatment should be distinguished from contemporary dermatological and psychiatric evidence the former does not establish the scientific validity or efficacy of homoeopathic treatment for psychodermatological disorders.

The importance of considering the patient's mental and emotional state alongside physical symptoms when constructing an individualised disease picture. This concept can be correlated with psychodermatology, where conditions such as eczema, psoriasis, chronic pruritus, and urticaria may be associated with psychological stress, anxiety, embarrassment, or emotional distress.[5] In accordance with homoeopathy, the patient's disposition may be documented as part of the clinical history, while §213–217 signifies the importance of assessing both physical manifestations and psychological experiences. The value of understanding the chronology of symptoms and integrating previous disease history with the present clinical picture. In a psychodermatological view, this approach involves documenting skin lesions, symptom severity, itching, sleep disturbance, stressful life events, coping mechanisms, and quality of life. However, Hahnemann's concepts of miasms and homoeopathic remedy selection should be distinguished from contemporary scientific explanations of mind–skin interactions.

§221–230 extend this discussion to acute and chronic mental disturbances, emotional influences on physical health, and the importance of appropriate patient management. These concepts can be correlated with psychodermatological conditions in which psychological stress may aggravate skin symptoms, while chronic or visible skin lesions may contribute to anxiety, depression, low self-esteem, and social withdrawal. Emotional factors as contributing to physical illness, which can be compared with contemporary observations that stress may influence itching and inflammatory skin conditions through interactions between the nervous, endocrine, and immune systems. §226, §228, and §229 emphasise supportive approaches and appropriate behaviour towards patients, which correspond to the modern principles of empathetic communication, psychological support, and non-judgmental care. Section 224 also signifies the importance of distinguishing mental illness from difficulties attributed to life circumstances, a distinction that requires careful clinical and psychological assessment today. Individualised homoeopathic treatment for mental and emotional disorders, its claims of therapeutic superiority are not established by contemporary evidence. Thus, the broader relevance of § 211–230 to psychodermatology lies in their emphasis on holistic case assessment, the relationship between mental and physical manifestations, and compassionate patient care, while treatment decisions should be guided by current dermatological and mental health evidence.

Relevance of Psychodermatology in Homoeopathic Case-Taking

Psychosomatic diseases can be viewed as the result of a complex interaction between psychological, biological, genetic and environmental factors. Emotional experiences such as stress, anxiety, grief, fear and psychological trauma may influence physiological functioning through neuroendocrine, autonomic nervous system and immune mechanisms. Individual susceptibility, personality traits, genetic predisposition and environmental influences further contribute to the development and progression of physical illnesses. Thus, the psychosomatic perspective emphasizes the interconnectedness of mind and body, signifying the importance of understanding an individual's emotional experiences, physical manifestations and personal history in comprehensive clinical assessment.[6]

Correlation through representative psychodermatological view

1. **Vitiligo:** The Interplay Between Psychological Stress and Autoimmune Mechanisms

Vitiligo is primarily an autoimmune-associated disorder in which autoreactive CD8+ T lymphocytes target melanocytes, leading to their destruction and loss of skin pigmentation.

Interferon-gamma signalling and inflammatory chemokines, particularly CXCL9 and CXCL10, contribute to the recruitment of immune cells involved in this process. The onset or progression of depigmented patches may follow stressful life events, emotional trauma or prolonged anxiety.[7]

2. Eczema associated with psychological stress

Eczema is a chronic inflammatory skin disorder characterised by dryness, itching, erythema and recurrent eczematous lesions. Its pathogenesis involves epidermal barrier dysfunction, genetic susceptibility, immune dysregulation and environmental influences. Psychological stress may act as an aggravating factor by influencing neuroimmune interactions and inflammatory responses.[8] Assessment should consider the relationship between emotional stress, clinical exacerbations and psychological burden. While stress may aggravate eczema, persistent itching and inflammation may further increase emotional distress. Clinical severity and quality of life can be evaluated using EASI and DLQI, respectively.[9]

3. Psoriasis with anxiety and social withdrawal

Psoriasis is a chronic, immune-mediated inflammatory skin disorder characterized by erythematous plaques covered with silvery-white scales. Stress-related neuroendocrine and immunological changes may influence inflammatory pathways involved in psoriasis, particularly those associated with the interleukin-23 (IL-23) and T-helper 17 (Th17) axis.[10] Patients experiencing prolonged emotional stress may report worsening of existing lesions or recurrent exacerbations.

DISCUSSION

The importance of considering local affections as part of the patient's overall clinical presentation rather than viewing them in isolation. This perspective can be correlated with contemporary psychodermatology, which recognises the bidirectional relationship between psychological wellbeing and skin disorders. Conditions such as eczema, psoriasis, chronic pruritus, and vitiligo may be associated with psychological distress, while the visibility, chronicity, and symptoms of these conditions can adversely affect emotional wellbeing and quality of life.

In psychodermatological practice, the significance of these concepts lies in the comprehensive assessment of the patient, including the nature and duration of skin lesions, symptom chronology, psychological stressors, sleep disturbance, coping mechanisms, and social functioning. For example, stress may aggravate itching in eczema, while visible psoriatic

lesions or vitiligo-associated depigmentation may contribute to anxiety, embarrassment, and social withdrawal. In vitiligo, psychological stress may potentially interact with neuroendocrine and immune pathways, but the disease is primarily associated with autoimmune destruction of melanocytes, and a direct causal relationship between stress and disease progression remains unproven. Similarly, the emphasis on mental and emotional symptoms in §211–230 can be related to the modern recognition that psychological and physical health are interconnected. Nevertheless, this correlation is conceptual rather than proof of scientific equivalence between Hahnemannian theory and contemporary psychoneuroimmunology. Therefore, the enduring clinical relevance of these sections lies in encouraging individualised history-taking, attention to the psychological burden of skin disease, empathetic communication, and a holistic understanding of patient experiences. Such an approach should complement, rather than replace, evidence-based dermatological treatment and appropriate psychological or psychiatric support. The integration of physical assessment and psychological care provides a more comprehensive approach to psychodermatological disorders, without assuming that emotional factors are the sole cause of skin disease or that homoeopathic treatment has established therapeutic efficacy. Hahnemann proposed that chronic diseases could be associated with underlying miasmatic influences, classified as psora, sycosis, and syphilis. Within this theoretical framework, miasmatic analysis is used to interpret the chronicity, recurrence, and individual manifestations of disease and to guide remedy selection.

CONCLUSION

Hahnemann provides the importance of mental and emotional symptoms, the chronology of physical and psychological manifestations, individualised assessment, and appropriate patient interaction. These themes can be correlated with modern psychodermatology, particularly in conditions such as atopic dermatitis, psoriasis, chronic pruritus, and vitiligo. The recognition that skin disorders and psychological wellbeing may influence one another. This supports a comprehensive, non-stigmatising approach that combines dermatological assessment with psychological evaluation when indicated. However, Hahnemann’s miasmatic explanations and claims for homoeopathic treatment should remain distinguished from established evidence on psychodermatological mechanisms and therapies.

REFERENCE

1. Pawlikowska-Łagód K, Polak A, Grzybowski A. Psychodermatology: From historic

concepts to contemporary clinical practice. Clin Dermatol. 2026 Mar-Apr;44(2):299-304. doi: 10.1016/j.clindermatol.2026.01.011. Epub 2026 Feb 2. PMID: 41638322.

2. Tuckman A. The Potential Psychological Impact of Skin Conditions. Dermatol Ther (Heidelb). 2017 Jan;7(Suppl 1):53-57. doi: 10.1007/s13555-016-0169-7. Epub 2017 Feb 1. PMID: 28150109; PMCID: PMC5289120.
3. Vidal Yucha SE, Tamamoto KA, Kaplan DL. The importance of the neuro-immuno-cutaneous system on human skin equivalent design. Cell Prolif. 2019 Nov;52(6):e12677. doi: 10.1111/cpr.12677. Epub 2019 Aug 23. PMID: 31441145; PMCID: PMC6869210.
4. Kowalewska B, Jankowiak B, Krajewska-Kułak E, Khvorik DF, Niczyporuk W. Quality of life in skin diseases as perceived by patients and nurses. Postepy Dermatol Alergol. 2020 Dec;37(6):956-961. doi: 10.5114/ada.2019.86182. Epub 2019 Jul 29. PMID: 33603616; PMCID: PMC7874861.
5. Jafferany M. Psychodermatology: a guide to understanding common psychocutaneous disorders. Prim Care Companion J Clin Psychiatry. 2007;9(3):203-13. doi: 10.4088/pcc.v09n0306. PMID: 17632653; PMCID: PMC1911167.
6. Tatayeva R, Ossadchaya E, Sarculova S, Sembayeva Z, Koigeldinova S. Psychosomatic Aspects of The Development of Comorbid Pathology: A Review. Med J Islam Repub Iran. 2022 Dec 12;36:152. doi: 10.47176/mjiri.36.152. PMID: 36636258; PMCID: PMC9826781.
7. Chang WL, Lee WR, Kuo YC, Huang YH. Vitiligo: An Autoimmune Skin Disease and its Immunomodulatory Therapeutic Intervention. Front Cell Dev Biol. 2021 Dec 14;9:797026. doi: 10.3389/fcell.2021.797026. PMID: 34970551; PMCID: PMC8712646.
8. Nemeth V, Syed HA, Evans J. Eczema. [Updated 2024 Mar 1]. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 Jan-. Available from: https://www.ncbi.nlm.nih.gov/books/NBK538209
9. Lönndahl L, Abdelhadi S, Holst M, Lonne-Rahm SB, Nordlind K, Johansson B. Psychological Stress and Atopic Dermatitis: A Focus Group Study. Ann Dermatol. 2023 Oct;35(5):342-347. doi: 10.5021/ad.22.035. PMID: 37830415; PMCID: PMC10579569.
10. Man AM, Orăsan MS, Hoteiuc OA, Olănescu-Vaida-Voevod MC, Mocan T. Inflammation and Psoriasis: A Comprehensive Review. Int J Mol Sci. 2023 Nov 8;24(22):16095. doi: 10.3390/ijms242216095. PMID: 38003284; PMCID: PMC10671208.