
COMPLICATIONS OF DIABETES MELLITUS

**Boda Shirisha, Bojja Srikarunya, Jakkula Sowmya, Mallarapu Akanksha, Kothuri Ashwini.*

Vaageswari College of Pharmacy, Karimnagar, India.

Article Received: 28 January 2026

*Corresponding Author: Boda Shirisha

Article Revised: 17 February 2026

Vaageswari College of Pharmacy, Karimnagar, India.

Published on: 09 March 2026

DOI: <https://doi-doi.org/101555/ijrpa.7930>

ABSTRACT:

Diabetes Mellitus, which is described through persistent hyperglycaemia that ultimately results in multi-organ failure, is a leading international health burden. Current knowledge of diabetes complications, their pathophysiological mechanisms, and their therapeutic implications is summarised in this review. Through oxidative stress, the manufacturing of advanced glycation end products, and inflammatory pathways, persistent hyperglycemia sets in a sequence of metabolic and vascular disruptions. Microvascular sequelae consist of retinopathy, nephropathy, and Neuropathy, whereas important microvascular and macrovascular problems include peripheral artery disorder, cerebrovascular accidents, and cardiovascular ailments. About 38 million Americans have diabetes, in keeping with current epidemiological information, and 1 in 5 go untreated, which will extensively increase morbidity and mortality. Between 50-80% of deaths from diabetes are because of cardiovascular issues, which stay the main motive of death. Diabetic nephropathy impacts 20-40% of patients, representing the primary cause of end-stage renal disorder in advanced international locations. About one-0.33 of patients have diabetic retinopathy, which is still the most common preventable cause of blindness in people of working age. Up to 50% of diabetes patients have peripheral Neuropathy, which dramatically raises the risk of foot ulcers and the need for amputation. With the addition of newly developed focused treatments, modern-day treatment approaches prioritise chance thing control, in depth glyceimic management, and early identification. Developing successful preventative measures and improving metabolisedults in diabetes treatment requires information on these outcomes.

KEYWORDS: Diabetes mellitus, microvascular headaches, macrovascular diabetic nephropathy, diabetic retinopathy, cardiovascular disorder, Neuropathy.

INTRODUCTION:

A metabolic situation referred to as persistent hyperglycemia may be caused by insufficient insulin secretion, poor insulin action both. Particularly, insulin is a crucial metabolic hormone affects how proteins, fat, and carbohydrates are metabolised. Because of insulin resistance, the metabolic abnormalities linked to diabetes generally affect tissues like skeletal muscle, adipose tissue, and the liver. The type and period of diabetes can affect how excessive the symptoms are. People who have high blood sugar levels, especially youngsters who don't have any insulin at all, may also have signs and symptoms like weight loss, accelerated urge for food, polydipsia, dysuria, and visual troubles. Some diabetic patients, particularly people with type 2 diabetes in its early stages, might not exhibit any signs and symptoms. Uncontrolled diabetes can cause several issues if left untreated, which include unconsciousness, disorientation, and, in rare cases, death from untreated nonketotic hyperosmolar syndrome or ketoacidosis.

According to a 2014 WHO document, eight.5% of humans over the age of 18 had diabetes. In 2019, diabetes prompted 1.5 million fatalities, forty eight% of which occurred in those under the age of 70. Roughly 20% of deaths from cardiovascular disease were related to high blood glucose, and diabetes also triggered a further 460,000 deaths from kidney disease. Standardised mortality rates from diabetes accelerated via 3% among 2000 and 2019. The diabetes-related mortality rate rose by thirteen per cent% in middle-income nations. However, between 2000 and 2019, the worldwide risk of death from any of the 4 fundamental non-communicable diseases (most cancers, diabetes, chronic respiratory diseases, or cardiovascular diseases) decreased through 22% for the ones elderly 30–70.

Patients with type 1 or type 2 diabetes regularly experience diabetes headaches, which may contribute substantially to morbidity and mortality. The major categories of diabetes persistent results are microvascular and macrovascular, with the previous being greater common than the latter. Neuropathy, nephropathy, and retinopathy are examples of microvascular issues, whereas peripheral artery disease, cardiovascular ailments, and stroke are examples of macrovascular issues. One of the primary causes of lower limb amputations is diabetic foot syndrome, that's characterized by the presence of foot ulcers related to Neuropathy, peripheral artery disease, and contamination. Lastly, there are extra diabetic issues that don't fall under the 2 formerly referred to categories, including dental ailments,

decreased infection resistance, and delivery headaches amongst ladies with gestational diabetes.

COMPLICATIONS:

Microvascular Complications:

Indices of Subclinical inflammation, along with better hsCRP, are correlated with the superiority of kind 2 diabetes and metabolic syndrome. Meprins are metalloproteinases expressed inside the Kidney by way of proximal tubules-that has been verified to play a great function in the development of diabetic nephropathy. In the paper titled “Meprin Metalloprotease Deficiency Associated with better mortality Rates & More intense Diabetic kidney Injury in Mice With STZ-Induced Type 1 Diabetes, J.E-Bylander et al. Designed experiments to evaluate the outcomes of meprin deficiency in mice with STZ-precipitated diabetes. Using meprin $\alpha\beta$ knockout (ko) mice, the authors tested the role of meprins in the improvement of diabetic nephropathy.

Monofilament & vibration notion checks are commonly used screening tools encouraged by means of several medical pointers to detect diabetic peripheral Neuropathy (DPN) or to assess the chance of foot ulcer formation. Monofilament checking out used as a screening tool for detecting DPN had limited sensitivity, but proper specificity. Dealing with the equal trouble, J.J. Brown et al., in any other paper of this unique trouble titled “An evaluation of Tingeening tools for the early detection of peripheral Neuropathy in adults with and without Type 2 Diabetes” InvesInvestigated the effectiveness of several tests for the early detection of DPN in patients with prediabetes or type 2 Diabetes.

Macrovascular Complications:

One most investigated comorbidities of diabetes is peripheral artery disorder. About 50% of individuals with diabetic foot ulcers also have PAD, and they’ll experience continual Ischemic soreness. Pain management can substantially improve quality of life for those individuals A. Tedeschi et al. Assessed the effectiveness and tolerability of Tapentadol PR, A medicine that has capabilities as a μ -opioid receptor agonist and a norepinephrine reuptake inhibitor, in patients with type 2 diabetes and severe chronic ischemic pain in a paper titled “Tapentadol prolonged release (PR) reduces the Severe chronic ischemic pain and l improves quality of life in patients with type 2 diabetes”. A numerical rating Scale (NRS) and Questionnaires, including DN4 and SF-12 Health Survey, were used to assess the analgesic efficacy of the drug. The outcomes of these drug studies guide the tapenta PR, which is an

efficient therapy for the treatment of chronic ischemic pain in sufferers with DM. The authors concluded that tapentadol PR substantially reduces pain intensity, Alleviates neuropathic signs, and complements the affected person's quality of lifestyle.

Miscellaneous Complications:

It has been shown that people with diabetes who are afflicted by severe hypoglycemic episodes are more likely to develop cardiovascular disease. A retrospective chart assessment study was conducted by S. Malkani and A. Kotwal using the university's digital database. In their paper titled "Blocking AGE-RAGE Signalling Improves Functional Disorders of Macrophages in the Diabetic Wound," Q. Wang et al. examined the function of RAGE-expressing macrophages in the failure of wound restoration in diabetic mice. There is accumulating evidence that the formation of advanced glycation end products (AGEs) may play an essential role in the impaired wound healing observed in patients with diabetes. Therefore, the researchers focused their attention on the wound-healing process. By topically administering anti-RAGE antibodies, the scientists attempted to inhibit AGE-RAGE signaling in the wound area. They observed that the group of mice that received anti-RAGE antibody treatment showed a significant acceleration in wound healing.

TREATMENT

Because hyperglycemia can change the lens of the eye in numerous ways, patients with type 2 diabetes are more likely to develop cataracts. The study titled "Diosgenin, a Novel Aldose Reductase Inhibitor, Attenuates the Galactosemic Cataract in Rats" by L. Ji et al. investigated the possible protective effect of diosgenin, an aldose reductase inhibitor (ARI), against the development of sugar-induced cataracts in rats.

The authors conducted two separate experiments. In the first experiment, rats were administered diosgenin, and it was observed that this AR inhibitor delayed the development of diabetic cataracts by delaying the opacification of the lens. In the second experiment, lens epithelial cells (LECs) derived from a cell culture were treated with diosgenin. The researchers found that the osmotic swelling of LECs decreased, which is a critical step in the onset and progression of diabetic cataracts. The authors' findings support the hypothesis that aldose reductase inhibitors can delay the onset of sugar cataracts.

Finally, S. Schacht et al. tested the effect of genistein, a naturally occurring phytoestrogen found in soy, on diabetes-related gastrointestinal disorders in mice in the article titled

“Dietary Genistein Influences Number of Acetylcholine Receptors in Female Diabetic Jejunum.” Compared with the control group, the intestinal contractions of diabetic mice were more widely spaced, and their jejunum contained fewer acetylcholine receptors (AChRs). A four-week genistein diet significantly restored the number of AChRs in the jejunum, although it did not affect the time interval between intestinal contractions. These results are encouraging because they suggest that genistein may be useful in treating gastrointestinal complications associated with diabetes.

REFERENCE:

1. Antar SA, Ashour NA, Sharaky M, Khattab M, Ashour NA, Zaid RT, Roh EJ, Elkamhawy A. Diabetes: Classification, mediators, and complications. *Biomedicine & Pharmacotherapy*. 2023 Dec;168:115734. Doi:10.1016/j.biopha.2023.115734.
2. Papatheodorou K, Banach M, Bekiari E, Rizzo M, Edmonds M. Complications of diabetes 2017. *Journal of Diabetes Research*. 2018 Mar 11;2018:3086167. Doi:10.1155/2018/3086167.
3. A.N. Orekhov, The diabetes mellitus–atherosclerosis connection: The role of lipid and glucose metabolism and chronic inflammation, *Int. J. Mol. Sci.*, 21 (5) (2020), p. 1835, 10.3390/ijms21051835
4. Abner EL, et al. Diabetes is associated with cerebrovascular but not Alzheimer’s disease neuropathology. *Alzheimer’s Dement.* 2016;12:882–889. doi: 10.1016/j.jalz.2015.12.006. [DOI] [PMC free article] [PubMed] [Google Scholar]
5. Langa KM, Levine DA. The diagnosis and management of mild cognitive impairment: a clinical review. *JAMA*. 2014;312:2551–2561. Doi: 10.1001/jama.2014.13806.